

Annual Report 2011

To order the 2011 annual report please contact:

**Deutsche Forschungsanstalt für
Lebensmittelchemie**

Mrs. A. Stoiber

Lise-Meitner-Strasse 34

D-85354 Freising, Germany

Phone: +49 8161 712928

Fax: +49 8161 712970

Anneliese.Stoiber@lrz.tum.de



Table of Contents

Structure and Bioactivity of Low-Molecular Food Ingredients (Hedonic Value)

- On the Formation of Methional during Pineapple Juice Processing
- Characterisation of Key Aroma Compounds in Dry Sausage
- Characterisation of Key Aroma Compounds in the Ginger Rhizome
- Characterisation of the Key Aroma Compounds in Dornfelder Red Wine
- Generation of Aroma-Active Fatty Acid Decomposition Products as well as Toxicologically Relevant Compounds During Deep-Frying of Potatoes
- Quantitative Study on the Aroma-Active Compounds of Cold-Pressed Rapeseed Oil
- Sunstruck Flavour Formation in Beer: Key Odorants and its Precursors
- Identification of Key Aroma Compounds in Wheat Beer
- Identification of the Key Aroma Compounds in Bartlett (Williams Christ) Pear Brandy
- Comparison of the Key Aroma Compounds in Raw and Roasted Styrian Pumpkin Seeds
- On the Contribution of 1-Alkene-1-Thiols to the Aroma of Roasted Sesame
- Structure-Odour Relationship in Homologous Series of Mercaptoalkyl Esters
- Quantitation of Key Odorants in Rape and Chestnut Honey
- Characterisation of the Key Aroma Compounds in Tomato Paste

Development of Analytical Methods

- Development of a Method to Quantify Common Wheat in Spelt Products
- Development of Stable Isotope Dilution Assays for the Quantitation of Glycidyl Esters in Edible Fats and Oils

Correlation between the Chemical Structure and the Physical Properties of Biopolymers

- Texture-Dependent Release of Sodium from Wheat Dough and Bread
- Structural Changes of Disulphide Bonds of Gluten Proteins during Bread Making
- Properties of Wheat Gluten as Affected by High-Pressure Induced Deamidation
- Comparison of Different Protein References and ELISA Kits for the Detection of Gluten in Foods

- Comparison of Methods for Quantifying the Glutenin Macropolymer Content of Wheat Flour
- Comparison of the Protein Composition of Different Wheat Species
- Gluten-Specific Peptidase Activity of Different Cereals as Affected by Germination
- Quantitative Protein Composition of Different Oat Cultivars
- Properties of Fish Filet as Affected by High-Pressure Treatment

Physiological Effects of Food Ingredients

- Trace Amine-Associated Receptor 1 (TAAR1) is Functionally Expressed in Human Leukocytes
- A New Cell System for the De-orphaning and Characterization of Trace-Amine-Associated Receptors
- Impact of [6]-, [8]- and [10]-Gingerol on Cytokine Secretion by Activated Human T-Lymphocytes
- Impact of Resveratrol Metabolites on the Cytotoxic Activity of Natural Killer Cells
- Effects of Catechol and Pyrogallol on CaCo-2 Cells: Elucidation of Structure Related Bioactivity via Impedance Based Cell Analysis