

## Annual Report 2012

To order the 2012 annual report please contact:

**Deutsche Forschungsanstalt für  
Lebensmittelchemie**

**Mrs. A. Stoiber**

**Lise-Meitner-Strasse 34**

**D-85354 Freising, Germany**

**Phone: +49 8161 712928**

**Fax: +49 8161 712970**

**[Anneliese.Stoiber@lrz.tum.de](mailto:Anneliese.Stoiber@lrz.tum.de)**



## Table of Contents

### Structure and Bioactivity of Low-Molecular Food Ingredients (Hedonic Value)

- Studies on the Formation Pathway of Acrolein
- Quantitation of Key Aroma Compounds in Raw Ginger
- Influence of the Manufacturing Process on Changes in Aroma Compounds of Dornfelder Red Wine
- Characterisation of Key Aroma Compounds in Wheat Beer
- Structural Elucidation of Aroma-Active Compounds in Durian
- Characterisation of the Key Aroma Compounds in Bartlett (Williams Christ) Pear Brandy
- Changes in the Odour-Active Compounds of French Oak Wood (*Quercus Robur*) During Toasting
- Characterisation of Important Aroma Compounds in Raw Liquorice (*Glycyrrhiza glabra*)
- Improvement of the Aroma of the Gluten-Free Bread by Addition of Corn Malt

### Development of Analytical Methods

- Profiling the Volatile Metabolome of Raw and Processed Hazelnuts by Comprehensive Two-Dimensional Gaschromatography (GC×GC/MS) and Novel Data Analysis
- Quantification of the Wheat Content of Spelt Products by a Stable Isotope Dilution Assay

### Correlation between the Chemical Structure and the Physical Properties of Biopolymers

- Salt Perception in Wheat Bread as Affected by the Distribution of Salt in the Crumb
- Generation of Malt with High Peptidase Activity for the Production of Gluten-Free Beer
- Collaborative Study on the Quantification of Gluten by R5 Sandwich-ELISA
- Quantification of Gluten Protein Fractions by Spectrophotometric Methods
- Functional Characterisation of Wheat Cultivars by a Gluten Aggregation Test
- Correlations Between Indirect Quality Parameters and the Baking Quality of Wheat Flours
- Reassessment of the Electrophoretic Mobility of High-Molecular-Weight Glutenin Subunits on SDS-Gels

- Biochemical Characterisation of Oat Proteins from the Globulin and Glutelin Fractions
- Solubility of Structure-Forming Proteins of Fish Muscle as Affected by High Pressure Treatment

### **Physiological Effects of Food Ingredients**

- Biogenic Amines Activate Immune Cell Functions via TAAR-Family Amine Receptors 1 and 2
- Quantification of Gingerols in Human Plasma via Stable Isotope Dilution Assay
- Odorant Receptor Amino Acid Motifs Govern the Functional Expression of Transmembrane Receptors

### **Food Composition Tables**

- Mineral Content of Different Teas