

Annual Report 2014

To order the 2014 annual report please contact:

**Deutsche Forschungsanstalt für Lebensmittelchemie
Leibniz Institut
Mrs. A. Stoiber
Lise-Meitner-Strasse 34
D-85354 Freising, Germany
Phone: +49 8161 712928
Fax: +49 8161 712970
Anneliese.Stoiber@lrz.tum.de**



Table of Contents

Structure and Bioactivity of Low-Molecular Food Ingredients (Hedonic Value)

- Influence of a Pasteurisation on Changes in the Aroma Compounds of Hand-Squeezed Oranges from Valencia Late
- Additive Effects of Esters on the Overall Aroma Quality of Apple Juices – Identification of Aroma-Relevant Esters as Markers for Juice Quality
- Characterisation of Key Aroma Compounds in Swiss Emmentaler Cheese
- Aroma-Active Compounds in Mangoes of Different Cultivars
- Characterisation of Important Aroma Compounds in Fresh Rosemary
- Influence of Curing on Selected Aroma Compounds and Biogenic Amines in Fermented Sausage
- Influence of Protein and Fat Content on the Aroma Release from a Milk Protein-Polysaccharide Foam Model
- Distribution of Selected Aroma Compounds Between the Liquid and Foam Phase in Beers with Different Foam Stabilities
- Regulation of the Flavour Formation in Wheat Pastries by Optimised Dough Proofing
- Amadori-Rearrangement Products as Possible Precursors of Strecker-Reaction-Products in Different Thermally Processed Foods

Development of Analytical Methods

- 3-MCPD-, 2-MCPD- and Glycidyl Fatty Acid Esters – Analysis, Stability and Reactivity
- Non-Immunochemical Gluten Quantitation in Wheat Starch: Pitfalls and Benefits
- Comparison of ELISA-Kits for the Quantitation of Gluten in Cereal-Based Foods
- Development of Stable Isotope Dilution Assays for the Quantitation of Aroma-Active 1,1-Dithio Compounds in Durian

Correlation between the Chemical Structure and the Physical Properties of Biopolymers

- Optimisation of Micro-Scale Extension Tests of Wheat Dough
- Baking Test to Determine the Functional Effects of Lipase-Treated Wheat Lipids
- Studies on the Effects of Transglutaminase and Xylanase in Rye Dough

- Colorimetric Quantitation of Quality-Related Gluten Protein Fractions Using Naphtol Blue Black
- Quantitative Composition of Osborne-Fractions of Wheat Flour as Affected by Blocking of Thiol Groups
- Factors Affecting the Isolation of Glutenin Macropolymer from Wheat Flour
- Protein Composition of Transgenic Winter Wheat after Ectopic Expression of a Barley Sucrose Transporter
- Influence of Artificial Drying of Maturing Wheat Grains on the Quantitative Protein Composition
- Salt Release and Distribution: New Ways of Reducing Sodium in Frozen Pizza
- Production of Gluten-Free Beer by Using Malt Extract with High Peptidase Activity
- Enzymatic Degradation of Gluten in Wheat Bran and Bread Drink
- Studies on the Synthesis and Detection of a 33-Mer Peptide from α -Gliadin

Physiological Effects of Food Ingredients

- The Human Carvone Receptor OR1A1 – from Enantiomer Specificity to Enantiomer Selectivity
- Influence of Glucuronic Acid Conjugates of Gingeroles on the Cytokine Secretion in Human T-Lymphocytes
- The Odorant Receptor OR2M3 Specifically Detects Key Food Odorants out of the Homologous Series of 3-Mercapto-2-Methylalkan-1-Ols – Correlation of EC_{50} -Values *In-Vitro* with Odour Thresholds *In-Vivo*

Food Composition Tables

- Gluten Content of Different Cereals and Cereal Products